

## **Stock assessment of Blue Mackerel Pacific stock (2019)**

Responsible Institute: National Research Institute of Fisheries Science

### Summary

The stock biomass was estimated by cohort model considering abundance index. Although the biomass fluctuated between 254 to 378 thousand tons during 1995 to 2003, it largely increased over 700 thousand tons in 2009 and 2010 fishing season by extremely high recruitment of 2004 and 2009. However, it decreased after 2011 by no significant high recruitment after 2010, it was 124 thousand tons in 2018 fishing season. The SSB increased same as the stock biomass between 165 and 340 thousand tons during 2006 to 2014 fishing seasons, then it decreased 53 thousand tons in 2018. Fishing pressure (average of fishery coefficient at age) revealed relatively stay at same level except high levels of 1995 to 1997, 2002 and 2003 fishing season.

For this population stock, we propose the Ricker (RI) model of reproduction curve for the stock-recruitment relationship equation,  $SB_{msy}$  is estimated 158 thousand tons (Yukami et al. 2019a). Following the reference, SSB at 2018 is below  $SB_{msy}$ . The fishing pressure on the stock are above  $F_{msy}$  since 2014. The status of SSB was considered at “decreasing” level by the past five years trend between 2014 to 2018.


- Summary table of reference relating to MSY

References	Values
<b>Regarding MSY</b>	
SBmsy	158,000 tons
Fmsy	0yr, 1yr, 2yr, 3yr, 4yr above= 0.22, 0.31, 0.48, 0.87, 0.87
%SPR(Fmsy)	27%
MSY	105,000 tons
<b>SSB and Fishing pressure at 2018</b>	
SB2018	53,000 tons
F2018	0yr, 1yr, 2yr, 3yr, 4yr above= 0.28, 0.34, 0.48, 0.83, 0.83
%SPR(F2018)	24%
%SPR(F2016-F2018)	21%
<b>Ratio to MSY</b>	
SB2018/SBmsy	0.33
F2018/Fmsy	1.09

- S-R relationship assumption: Ricker (without autocorrelation)

- Summary of stock status

Status of current SSB	Below SBmsy
Status of F	Above Fmsy
Status of SSB	decreasing



The relationship between SB/SBmsy and F/Fmsy.

The values of three years moving average were used for both SB and fishing intensity.

## 1. Data set

The data set used for the stock assessment is as follows.

Data set	Data source and research
Catch number by age and year	Landing at major ports (Hokkaido-Miyazaki [17] prefectures, JAFIC, Northern Pulse Seine Fisheries Association) Length composition by month (NRIFS, Hokkaido-Miyazaki [17] prefectures, JAFIC); market measurement Length, weight, age, and maturity data by month (NRIFS, Hokkaido-Miyazaki [17] prefectures, JAFIC); market measurement, fisheries testing Mixing rate research of chub and blue mackerel by month and fisheries (NRIFS, Hokkaido-Miyazaki [17] Prefectures); at sample ports, market samples, and fisheries research
<b>Index of the stock</b>	
Recruitment index	East Hokkaido-Sanriku Waters drift-net fishing survey CPUE (Hokkaido); drift net Mid water trawl survey at Northwestern Pacific(May-July, NRIFS); mid water trawl* Mid water trawl survey at Northwestern Pacific (September/October, NRIFS); mid water trawl*
SSB index	Izu Islands waters dip-net fishing CPUE (Kanagawa Prefecture) *
Spawning volume	Egg and larvae survey (NRIFS, Aomori-Miyazaki [18] prefectures); NORPAC net
Natural mortality	Assuming $M = 0.4$ per year (Tanaka 1960)

\*: used as the tuning index for the cohort analysis.

## 2. Ecology of the species

### 1) Distribution and migration

Blue mackerel tends to distribute warm offshore waters (Ochiai and Tanaka 1998). The main distribution area for adults is around water of the Kuroshio current.

The distribution and migration are shown in Fig. 2-1. The larvae hatched around the Kuroshio current and distributes from the coastal water of southern Honsyu to the transition water between Kuroshio and Oyashio currents at located 165 to 170 East longitude as same as the chub mackerel larvae (Watanabe et al. 1999, Nishida et al. 2000, Kawabata et al. 2006a). The juveniles sized at 5 to 15cm FL transferred to transition water, migrate to north with growing, feed at the area from coastal water of eastern Hakkaido and Krill Islands to the subarctic water around 165 degree East longitude where the surface temperature around 13°C in summer to fall (Savinykh et al. 2004, Kawabata et al. 2006a and 2007). It reached 20 to 25cm FL in fall to winter, and down south to the coastal waters of Joban and Boso to offshore water around Kuroshio current for wintering (Kawabata et al. 2009b). The wintering ground at the water near Emperor Sea mounts was found for 2004 year class which had high recruitment (Kawabata et al. 2008 and 2009a). Age one fish did not appear the water

north of Sanriku district after wintering until 1980, but it became migrating to the water from Tohoku to Hokkaido according to the increase of surface temperature since 2001 (Kawabata et al. 2006b and 2008). It down south for wintering and migrate to the Izu Islands water for spawning in spring (Meguro et al. 2002). Many schools distribute near Kuroshio current at the coastal water of southern Honshu all the year to be targeted by many fisheries, which are different from the schools largely migrate from near the Kuroshio current at the Izu Island to Tohoku and Hokkaido waters. It is suggested that many fish above age three does not migrate north of Sanriku district and stay at the western water near the cape Ashizuri with small migrations or stay near the spawning grounds (Nashida et al. 2006). Furthermore, it is considered that the observation of schools mainly consisting with age 8 fish at the Emperor sea mount area in 2008 to 2015 were due to the dominant recruitment spawned at the water south of Hachijo Island (Kamimura et al 2017).

## 2) Age and growth

The larvae grow 1mm per day until 5cm FL after hatching observed by otolith reading (Watanabe et al. 2002), then it grows 15cm after 80days, and over 20cm of 120 days after hatching (Takahashi et al. 2010). The scale annuli reading is practical for the fish after subadult stage (Kondo and Kuroda 1966, Watanabe et al. 2002), it is used the survey. Otolith annuli and daily ring readings are also effective for age determination (Higuchi 1999, Kimura et al. 2002, Nashida et al. 2003, Katayama and Ishii 2009). Recent analysis for age and growth from sampling of catch indicates fish becoming 20-25cm FL at age 0 in fall, 28-31cm at age 1 in summer, 30-34cm at age 2, 33-36cm at age 3, around 37cm at age 4, and 45cm at the maximum. The longevity was estimated around age 6 from size composition of catch, but the oldest age 11 was reported. The growths at younger ages are different by area, and it in the western area of offshore Kumano tends to faster than fish occur in the water north of Izu Islands. The average length (FL), weight (average weight in catch in 2014 to 2018) by age used for future projection are shown in Fig. 2-2.

## 3) Reproduction

The blue mackerel mature and spawn above 30cm FL from the observation of ovary tissue (Hanai and Meguro 1997). The mature age was considered age 2 and above, then it is assumed that all the fish age 2 and above are mature and spawn (Fig. 2-2, 2-3). The spawning grounds are found from the waters southern Kyusyu and cape Ashizuri to the Kuroshio current water near Izu Islands (Tanoue 1966, Fig 2-1). The recruitments hatched at the larger spawning ground in the East china sea supposed to migrate into the Pacific waters (Tanoue 1966, Niiya 2007). A spawning season are from December to June next year at the western waters of cape Ashizuri, January to March in the East china sea, and February to March near the water of cape Ashizuri (Tanoue 1966, Nashida et al.

2006). The spawning season of main spawning ground of blue mackerel near Izu Island are March to June, but it is considered that it is not suitable as spawning grounds by the short spawning season from the ovary tissue observation and small amount of spawning eggs sampled (Watanabe et al. 2000, Hashimoto et al. 2005). However, it is supposed that larvae and juvenile occur in the north of transition area consist by the fish hatched at the Izu Island spawning grounds in March to June as same as chub mackerel (Takahashi et al. 2010).

#### 4) Prey-predator relationships

Larvae feed on planktonic crustaceans and larvae of anchovy or sardines (Ochiai and Tanaka 1998). After juvenile, it feed small teleost and cephalopods with preys mentioned above. It feeds fishes including anchovy, benttooth and lantern fishes, crustaceans like krill and cephalopods at the Kumano Nada fishing ground, horned krill and anchovy at Sanriku fishing ground and copepod, krill, anchovy, lantern fishes, cephalopod like Eupoloteuthidae and salpa in the transition area between Kuroshio and Oyashio where located offshore of Joban and Sanriku (Hotta 1957 and Yokota et al. 1961). Feeding by whales are observed at the high abundance periods (Matsuoka et al. 2008).

### 3. Fisheries on the species

#### 1) Outline of fisheries

The major fisheries for blue mackerel are middle scale purse seine, large scale purse seine, dip-net fishing and stick-held dip-net fishing (mid-area), set net and line fishing (Fig. 3-1). Fishing grounds are formed in the water on continental shelves and slopes, around water of Islands and slopes. Targeted fish are below age 2 and elder fishes larger than 40cm are rare in the catch. In the firelight mackerel fishing, main target are age 1 and 2 fishes. Line fishing operated at the southern waters targets adults staying at shallows, therefore it catches elder fishes. In the set net fishery, catches differ by area and season, and juveniles are caught in the southern area. The mixer catch with chub mackerels frequently occurs in the northern areas. The purse seine catch is the largest in various fisheries.

#### 2) History of catch

The blue mackerel catch was estimated from the mixing ratio survey of landing. Chub and blue mackerels are caught together by the fisheries and summed together as “mackerels” in fishery statistics. The identification of each species by external form are relatively easy which blue has clear black spots in side body, and fin base distance of first dorsal spine is narrower than chub mackerel (Fishery Agency 1999). Natural hybrid between blue and chub mackerels are found, but it is not considered a big issue

for stock assessment because occurrence rate is only 0.3% of total (Taniguchi et al. 1989 and Saito 2001).

For the assessment, fishing season, July to June next year was used as an unit of year, catch was summed by fishing year, and stock abundance was defined as the value of July. The catch by fishery fluctuated between one (1982) to 89 thousand tons (2006) for mid-scale purse seine, 4 (2018) to 62 thousand tons (1985) for firelight mackerel fishing, one (1991) to 32 thousand tons (2010) for set net, 0 (1994) to 64 thousand tons (2009) for northern purse seine, respectively (Fig. 3-1, Table 3-1). Especially for the northern purse seine, it decreased from 41 thousand tons in 2014 to 16 thousand in 2015, 3 thousand in 2016 and finally 490 tons in 2018. In total, catch increased high level from 100 thousand in 1995, to historical highest 193 thousand tons in 2006 by high recruitment of 2004. Thereafter, it remained high level at 191 thousand tons in 2010, then it decreased since 2011 and catch recorded at 34 thousand tons in 2018 as the lowest since 1995.

Before 1981, catch records as blue mackerel was not remained, but catch by the northern purse seine and set net were fairly small (Sou et al. 1980, TNFRI unpublished). Purse seine catch was small in the mid-area of Pacific coast, catch of dip net fishing mainly consisted with chub mackerel, and blue mackerel catch increased after the decreasing of chub mackerel since 1982 (Meguro 1999). Estimated blue mackerel catch was relatively smaller than that of after 1982 in the southern area of Pacific coast. It is supposed that the blue mackerel abundance before 1981 was largely smaller than that after 1995 which abundance was estimated (Fig. 3-1).

Recently, China and Russia catch chub mackerel at the high sea of North Pacific and 200 EEZ, respectively. They reported their catch after 2014 to the North Pacific Fisheries Commission (NPFC). Since the catch of both countries included blue and chub mackerel together, the blue mackerel catch by each country were estimated assuming the same mixture ratios of Japanese purse seine catch in July to December (Fig.3-1, Table 3-1). In the results, the ratios of blue mackerel were estimated at 21% in 2014, 9% in 2015, 2% in 2016, 1% in 2017, and 0.5% in 2018. The difference of statistical unit of catch, e.g. calendar year used by NPFC and fishing year used for analysis was supposed not an issue, because fishing season of both countries were July to November.

Age composition of catch mainly consist with age 1 and 2 fish with yearly large fluctuation due to the large contribution of purse seine fishery (Fig. 3-2, Appendix 3). When strong recruitment occurs, it appears in the catch as age 0 and 1 fish. The proportion of age 0 fish in the catch is relatively small except dominant year class. It is considered that the difference of the migration route and fishing grounds caused the less catchability of age 0 fish.

#### 4. Stock status

##### 1) Stock assessment methods

We estimated the numbers at age by years 1995-2018 using the tuning-VPA method (cohort analysis) which applies the Pope's approximation formula with taking fishing season (July to next June) as year and age 4 and above fish as age plus group (Pope, 1972) (Table 4-1, Appendices 1,2,3 and 4). The natural mortality coefficient ( $M$ ) was set at 0.4 per year estimated by the statistical relation between longevity and natural mortality,  $M=2.5/\text{longevity}$ , here longevity is 6 years then  $M$  became 0.4 (Equation of Tauchi and Tanaka, Tanaka 1960). Japanese catch at age was made by summing up catch at age of each prefecture where converted from catch at size calculated by length composition and length-weight relation of each three month of landing at major fishing port of Pacific coast. The catch at age of Russia and China were obtained assuming same as it of Japanese one came from north of Ensyu Nada area in July to December. Terminal  $F$  (2018) was determined with recruitment index obtained from stock held dip net fishery at Shizuoka prefecture (Fig. 4-1, Table 4-2) and SB index from standardized amount of spawning eggs (Appendix 3, Fig.4-2, Table 4-2) using as tuning index.

##### 2) Changes in the biomass indices

The yearly fluctuation of recruitment index is shown in Fig. 4-1 and Table 4-2. All indices are revealing same trends of recruitment level with high abundance in 1996, 2004 and 2009, and low level in 2006 and 2008. However, the results of cohort analysis do not indicate high level of recruitment in 2013, 2016, 2017 and 2018, but indices of northern migration and survey in fall indicated high recruitment. The assessment results and indices tend to be not fit recently.

The yearly fluctuations of amounts of eggs spawn as index of adults are shown in Fig. 4-2 and Table 4-2. After 2005, it indicated high values in 2007 and 2011 reflecting increase of mature fish in 2004 and 2009. Thereafter, it stayed around 25 trillion eggs until 2014, then it decreased to 8.9 trillion eggs. It increased again to 109.6 trillion eggs in 2018 then suddenly dropped to 10.9 trillion eggs in 2019 (only January to June). The 102.2 trillion eggs of 2018 total eggs suggested to be spawn at Area III (Hyuga Nada, Tosa bay, Cape Shionomisaki). It is considered that the total egg amount was over estimated considering frequency distribution of egg diameter, and fishing status of adult fish (Yukami et al. 2019b). It is considered that spawning condition in 2019 is similar of 2018, then concluded that the value of total eggs spawn cannot be used as tuning index, then the values are standardized (Appendix 3). The standardized values of amount eggs spawn peaked at 2008, then decreased to the lowest value in 2019 since 2005 (Fig. 4-2, Table 4-2).

### 3) Trends in biomass and fishing pressure

Biomass increased from around 300 thousand tons in 1995 to 600 thousand tons in 2004 by high recruitments of 1996 and 2004, then up to 700 thousand tons in 2009 and 2010 with high recruitment of 2009 (Fig.4-3, Table 4-1, Appendix 4). After 2011, it decreased to 113 thousand tons in 2017, the lowest after 1995, then increased to 124 thousand tons in 2018. The SB revealed same trend of biomass, in high level of 165 to 340 thousand tons in 2006 to 2014, then decreased to 53 thousand tons in 2018 (Fig. 4-3, Table 4-1, Appendix 4). The recruitment per spawning (RPS) stayed in stable except high value of 2004 since 1996, it decreased and stay low level after 2010 (Fig. 4-4).

As a sensitivity test on  $M$ , we used 0.3 and 0.5 as alternatives to the base value of 0.4 and estimated the biomass, SSB and recruitment for the most recent year 2018 (Fig. 4-5). The biomass ranged from 80% to 128%, SSB from 83% to 122% and recruitment from 77% to 134% compared to the base value, resulting in higher value as  $M$  increases.

Average  $F$  among all ages (simple average of  $F$  at age) fluctuated among years at a range of 0.44 and 1.15, and show relatively stable status except the high values of 1995 to 1997 and 2002 and 2003 (Figure 4-6, Appendix 4). The fishing ratio fluctuated ranging 21 to 56%, but stayed around 30% except high values of 1995 to 1997 (Fig. 4-7, Table 4-1, Appendix 4).

Item	Value	Remarks
SB2018	53 thousand tons	SSB in 2018
F2018	(age 0, 1, 2, 3, 4+)=(0.28, 0.34, 0.48, 0.83, 0.83)	
U2018	28%	Fishing ratio in 2018



4) Yield per recruitment (YPR) , spawning per recruitment (SPR) and current fishing pressure

In order to compare the fishing pressure considering the influence of selectivity, Figure 4-8 shows the %SPR (ratio of SPR which assumes no fishing divided by the SPR with current catch) calculated by converting the  $F$  value of each year. The lower the fishing pressure, the higher the %SPR. The %SPR were low during 1995 to 1997, then stayed around 25% during 1998 to 2013, and it slightly decreased in 2014. It gradually increased since 2015.

Figure 4-9 shows the relationship between YPR and %SPR for the current fishing pressure. As for the selectivity in  $F$ , we use the selectivity which was used to estimate  $F$  that gives MSY ( $F_{msy}$ ) which was suggested at the ‘Research Institute meeting on Reference points’ held in April, 2019.  $F_{msy}$  corresponds to 27% in %SPR. Current fishing pressure (F2014-F2018) is higher than  $F_{msy}$  and F30%SPR.

Item	Value	Remarks
%SPR (F2018)	24%	%SPR in 2018
%SPR (F2016-2018)	21%	%SPR corresponding to current fishing pressure (F2016-F2018)

5) Stock-recruitment relationship

Figure 4-10 shows the Stock-recruitment (S-R) relationship between SSB (in biomass) and recruitment (in numbers). According to the ‘Research Institute meeting on Reference points’ mentioned above, it is suggested to use the Ricker type model for the S-R relationship of this stock (Yukami *et al.* 2019a). Parameters for the S-R relationship is estimated based on the SSB and recruitment which are estimated by the stock assessment conducted in 2018, and as for the optimization method, least-squares method is used. The model does not consider auto-correlation between the residuals of the recruitment. Estimated parameters for the S-R relationship are shown in the Table below.

S-R relationship	Optimization method	Auto-correlation	$a$	$b$	S.D.
Ricker (RI) type	Least square	No	13.5	0.00558	0.507

Here,  $a$  and  $b$  are parameters of Ricker type reproduction model shown below.  $R$  is recruitment (million fish) and  $E$  is SB (thousand tons).

$$R=aEe^{-bE}$$

- 6) Level of SSB and fishing pressure that will achieve MSY under the current condition.

The table below shows the SSB and  $F$  that will achieve MSY (SBmsy, Fmsy) under the current environmental condition since 1995, which was suggested at the ‘Research Institute meeting on Reference points’ (Yukami *et al.* 2019a).

Item	Suggested value	Remarks
SBmsy	158 thousand tons	SSB that will obtain MSY
Fmsy	(age 0, 1,2,3,4+)= (0.22, 0.31, 0.48, 0.87, 0.87)	
%SPR (Fmsy)	27%	%SPR corresponding to Fmsy
MSY	105 thousand tons	MSY

- 7) Stock status, stock trend and level of fishing pressure

Figure 4-11 and Appendix 5 shows a Kobe-plot which shows the relationship between SSB and fishing pressure.  $F/F_{msy}$  shows the yearly ratio between  $F$  and  $F_{msy}$  under the current selectivity that gives  $F_{msy}$  which was converted to %SPR. It is regarded that the fishing pressure of this stock has been above the level that gives MSY until 2006. It stayed around  $F_{msy}$  during 2007 to 2013, then it became above  $F_{msy}$  again after 2014, it gradually decreased but was 1.09 times of  $F_{msy}$  in 2018. Although, the SSB were above SBmsy during 2006 to 2014, it was below SBmsy except that period. The SSB in 2018 was 0.33 times of the SBmsy. The condition of SSB is considered “decreasing” by the five years trend of SB (2014 to 2018).

Item	Value	Remarks
SB2018/ SBmsy	0.33	Ratio between the SSB that gives MSY and the SSB in 2018
$F_{2018}/ F_{msy}$	1.09	Ratio between the fishing pressure that gives MSY and the fishing pressure in 2018 *

\* Ratio between  $F_{2018}$  and  $F$  under the current selectivity that gives  $F_{msy}$  which was converted to %SPR.

Level of SSB	below SBmsy
Level of $F$	above $F_{msy}$
Trends in SSB	decreasing

## 5. Stock assessment summary

Biomass fluctuated between 254 and 378 thousand tons during 1995 to 2003, it reached over 700 thousand level in 2009 and 2010 by the strong recruitments of 2004 and 2009. However, no strong recruitment occurred after 2010, the biomass rapidly down to 124 thousand tons in 2018. The SB showed similar trend of biomass, it stayed

between 165 to 340 thousand tons during 2006 to 2014, then decreased and down to 53 thousand tons in 2018. Fishing pressure (average of  $F$  at age) were relatively stable except high level during 1995 to 1997, and 2002 to 2003.

The SB2018 is below the level of SB<sub>msy</sub>.  $F$  has been higher than the  $F_{msy}$  since 2014. Trends in SSB is considered to be “decreasing” based on the recent five-year trends (2014-2018).

## 6. Other matters

It is considered that the effect of catch regulation on age 0 fish to increase abundance of stock might be small, because fishing pressure on age 0 fish is not high and is less reliable increasing recruitment by increasing SB. However, it is valuable to consider HCR by age in terms of economy effects because fish age one and above having higher price than age 0 fish.

The stock conditions are different between chub and blue mackerels, for instance dominant year class occurs in chub mackerel recently but not for blue mackerel. While the blue mackerel is managed in the same TAC of chub mackerel due to the mixed catch with chub. Therefore it is important that monitoring of the stock should be conducted with chub mackerel stocks.

## 7. References

- 花井孝之 (1999) 伊豆諸島海域におけるゴマサバの資源特性について. 中央ブロック長期漁海況予報, **107**, 32-39.
- 花井孝之・目黒清美 (1997) ゴマサバの卵巣組織観察による成熟, 産卵についての基礎的研究. 関東近海のマサバについて, **30**, 92-99.
- 橋本 浩・池上直也・森 訓由・岡部 久 (2005) 2005年の関東近海におけるサバ属卵分布. 2005年度水産海洋学会大会講演要旨集, 120.
- 堀田秀之 (1957) カツオの胃内容物中にみられたゴマサバの幼・稚魚 (薩南海区). 東北水研研報, **9**, 129-132.
- 飯塚景記 (1978) 東北海区北部海域におけるゴマサバについての二・三の生物学的観察. 東北水研研報, **39**, 11-20.
- 上村泰洋・川端 淳・米崎史郎・高橋正知・由上龍嗣・渡邊千夏子 (2017) 天皇海山海域におけるゴマサバの生物学的特性. 水産海洋研究, **81**, 18-28.
- 片山知史・石井光廣 (2009) サバ類の耳石による年齢査定を試み. 2009年度水産海洋学会研究発表大会講演要旨集, 89.
- 川端 淳・中神正康・巢山 哲・西田 宏・渡邊千夏子 (2007) 北西太平洋における近年のゴマサバ0歳魚の分布, 回遊と加入量. 2007年度水産海洋学会大会講演要旨集, 9.
- 川端 淳・中神正康・巢山 哲・西田 宏・渡邊千夏子 (2008) 北西太平洋における近年のゴマサバ資源の増加と1歳魚以上の分布, 回遊. 黒潮の資源海洋研究, **9**, 61-66.

川端 淳・中神正康・巢山 哲・上野康弘・谷津明彦 (2009a) 2001～2008年5～7月の北西太平洋におけるサバ類0歳魚の分布、体長組成と加入豊度との関係. 2009年度水産海洋学会大会講演要旨集, 19.

川端 淳・中神正康・巢山 哲・谷津明彦・高木香織・建田夕帆 (2006a) 最近の広域な調査船調査から推定される北西太平洋におけるサバ, イワシ類の季節的分布回遊. 2006年度水産海洋学会大会講演要旨集, 94.

川端 淳・山口閔常・巢山 哲・中神正康 (2006b) 近年の東北～北海道海域における表層性魚類相とゴマサバの来遊動向. 月刊海洋, **38**, 175-180.

川端 淳・谷津明彦・西田 宏・小澤竜太・高木香織・山下紀生・山下夕帆・中神正康・高橋正知 (2009b) 北西太平洋におけるマサバ・ゴマサバ未成魚の越冬海域の年変化. 第57回サンマ等小型浮魚資源研究会議報告, 東北区水産研究所八戸支所, 157-162.

木村 量・梨田一也・大関芳沖・本多 仁 (2002) ゴマサバ *Scomber australasicus* に適した耳石による年齢査定法. 水産海洋研究, **66**, 247-251.

近藤恵一・黒田一紀 (1966) サバ属魚類の成長—I. 東海水研報, **45**, 31-60.

Matsuoka, K., S. Otani, T. Isoda, A. Wada, S. Kumagai, T. Ohshima, I. Yoshimura, K. Sugiyama, M. Aki, K. Kato, M.M.U. Bhuiyan, N. Funasaka, Y. Suzuki, R. Sudo, Y. Motohashi, M. Mori, M. Tsunekawa, D. Inagake, H. Murase and T. Ogawa (2008) Cruise report of the second phase of the Japanese Whale Research Program under Special Permit in the Western North Pacific (JARPN II) in 2007 - Offshore component -. Paper SC/60/O5 submitted to the 60th IWC Scientific Committee, 40pp. (※著者らは本調査の胃内容物標本を日鯨研を通じて実際に観察し, さば類はマサバとゴマサバであることを確認した)

目黒清美 (1999) 関東近海のゴマサバの分布について. 中央ブロック長期漁海況予報, **107**, 40-54.

目黒清美・梨田一也・三谷卓美・西田 宏・川端 淳 (2002) マサバとゴマサバの分布と回遊—成魚. 月刊海洋, **34**, 256-260.

梨田一也・本多 仁・阪地英男・木村 量 (2003) 足摺岬周辺及び土佐湾中央部海域で漁獲されたゴマサバの年齢形質としての耳石の有効性. 黒潮の資源海洋研究, **4**, 5-9.

梨田一也・本多 仁・阪地英男・三谷卓美・平井一行・上原伸二 (2006) 足摺岬周辺海域及び伊豆諸島海域で実施した標識放流調査によるゴマサバの移動・回遊. 水研センター研報, **17**, 1-15.

新谷淑生 (2007) 高知県西部海域におけるゴマサバ若魚の加入について. 黒潮の資源海洋研究, **8**, 101.

西田 宏・渡邊千夏子・谷津明彦・木下貴裕 (2000) 黒潮統流～黒潮親潮移行域における幼稚魚採集と表面水温情報を利用したマサバ・ゴマサバの加入量予測. 関東近海のマサバについて, **33**, 96-102.

落合 明・田中 克 (1998) ゴマサバ. 新版魚類学 (下) 改訂版, 恒星社厚生閣, 東京, 844-855.

Pope, J.G. (1972) An investigation of the accuracy of virtual population analysis using cohort analysis. Int. Comm. Northwest Atl. Fish. Res. Bull., **9**, 65-74.

斉藤憲治 (2001) リボゾーム DNA の変異を利用した種判別法. 東北水研ニュース, **62**, 2-5.

Savinykh, V.F., A.A. Baitalyuk and A.Yu. Zhigalin (2004) Pelagic fish new to the Pacific waters of the Southern Kurils, migrants from the zone of Kuroshio. *Journal of Ichthyology (Voprosy Ikhtiologii)*, **44**, 611-615.

水産庁 (1999) マサバ・ゴマサバ判別マニュアル. 水産庁水産業関係試験研究推進会議マサバ・ゴマサバ判別マニュアル作成ワーキンググループ, 中央水産研究所, 32 pp.

高橋正知・高木香織・川端 淳・渡邊千夏子・西田 宏・山下紀生・森 賢・巢山 哲・中神正康・上野康弘・斉藤真美 (2010) マサバ・ゴマサバ太平洋系群 2007 年級群の推定孵化時期. *黒潮の資源海洋研究*, **11**, 49-54.

田中昌一 (1960) 水産生物の Population Dynamics と漁業資源管理. *東海水研報*, **28**, 1-200.

谷口順彦・向井龍男・関 伸吾・津田恭敬 (1989) マサバ・ゴマサバ. *アイソザイムによる魚介類の集団解析, 海洋生物集団の識別等に関する先導的評価手法の開発事業報告書*, 日本水産資源保護協会, 371-384.

Tanoue, T. (1966) Studies on the seasonal migration and reproduction of the spotted mackerel, *Pneumatophorus tapeinocephalus* (BLEEKER). *Memoir of Fac. Fish. Kagoshima Univ.*, **15**, 91-175.

樋田史郎 (1999) ゴマサバの日齢査定について. *中央ブロック長期漁海況予報*, **107**, 83-91.

曾 萬年・中田英昭・平野敏行 (1980) 近年のゴマサバ資源の増大について. *水産海洋研究会報*, **36**, 19-26.

由上龍嗣・井須小羊子・上村泰洋・古市 生 (2019a) 平成 31 (2019) 年度ゴマサバ太平洋系群の管理基準値等に関する研究機関会議報告書. [http://www.fra.affrc.go.jp/shigen\\_hyok\\_a/SCmeeting/2019-1/detail\\_gomasaba\\_p.pdf](http://www.fra.affrc.go.jp/shigen_hyok_a/SCmeeting/2019-1/detail_gomasaba_p.pdf) (last accessed 30 October 2019)

由上龍嗣・井須小羊子・渡邊千夏子・上村泰洋・古市 生 (2019b) 平成 30 (2018) 年度ゴマサバ太平洋系群の資源評価. 平成 30 年度我が国周辺水域の漁業資源評価 第 1 分冊, 水産庁・水産研究・教育機構, 248-278.

渡邊千夏子・花井孝之・目黒清美 (2000) マサバとゴマサバの産卵生態の比較. 一日当たり総産卵量に基づくマサバ太平洋系群の資源量推定法に関する調査報告書, 中央水産研究所, 14-23.

渡邊千夏子・川端 淳・和田時夫 (1999) 黒潮親潮移行域におけるサバ類当歳魚の分布. *月刊海洋*, **31**, 236-240.

渡邊千夏子・小林憲一・川端 淳・梨田一也 (2002) マサバとゴマサバの年齢と成長. *月刊海洋*, **34**, 261-265.

山川 卓 (1999) 熊野灘におけるゴマサバの漁獲状況と尾叉長組成. *中央ブロック長期漁海況予報*, **107**, 25-39.

横田滝雄・通山正弘・金井富久子・野村星二 (1961) 魚食性魚類の胃内容物の研究. *南海水研報*, **14**, 153-202.

(執筆者：由上龍嗣、井須小羊子、上村泰洋、古市 生、渡部亮介、金森由妃)




Figure 2-1. Distribution and migration of blue mackerel Pacific stock.




Figure 2-2. Relationship between age and length (fork length), and between age and weight. Black triangle shows that of weight and white circle shows that of length.








Figure 2-3. Maturity rate by age.




**Figure 3-1.** Annual catches of blue mackerel by fisheries. Calendar year only Japan (above: January to December) and Fishing year including foreign countries (below: July to next June). The values for China and Russia are estimated values.



**Figure 3-2.** Annual age composition in catch. (Red: above 4 years old, green: 3 years old, blue: 2 years old, yellow: 1 year old, white: age 0).




**Figure 4-1.** Fluctuations in the recruitment index in various surveys. The indices are relative values with the index maximum and minimum set at 100 and 0, respectively.




**Figure 4-2.** Spawning eggs of blue mackerels in the Pacific waters off Japan. Values for 2019 is for January to June. Left axis is the number of spawning eggs (in trillion) and right axis is the standardized spawning eggs. Black square shows the spawning eggs and white circle shows the standardized spawning eggs.







**Figure 4-3.** Fluctuations in biomass and SSB. The white dots show SSB, and the black dots show biomass.




**Figure 4-4.** Fluctuations in recruitment and recruitment per spawning (RPS). White bar plot shows recruitment and black dots indicate RPS. Left y-axis is recruitment (100 million in numbers), right y-axis is the RPS (numbers/kg).




**Figure 4-5.** Biomass, SSB and recruitment for 2018 depending on the value of  $M$ . Black circle represents the biomass, square represents SSB and triangle represents the recruitment. Left y-axis is the biomass and SSB (thousand tons), right y-axis is the recruitment (100 million in numbers).




**Figure 4-6.** Fluctuations in  $F$  at age. The black solid line shows the average of all ages.




**Figure 4-7.** Fluctuations in catch proportion. Y-axis is the catch proportion (%).




**Figure 4-8.** Fluctuations in %SPR by years. %SPR shows the ratio of SSB when no fishing to the SSB when there is fishing, and %SPR becomes low with high  $F$  and vice versa.



**Figure 4-9.** Relationship between the fishing mortality (F, simple average) and %SPR, YPR.



**Figure 4-10.** Relationship between SSB and recruitment. The blue line shows the S-R relationship suggested at the ‘Research Institute meeting on Reference points’ held in April 2019 (Yukami et al. 2019a).



**Figure 4-11.** Relationship between the SSB and F that gives MSY (SBmsy, Fmsy) to the current SSB and F (Kobe plot). The  $F$  and SSB is the three-year moving average. The value in 2018 is the average of 2016 to 2018.

**Table 3-1.** Annual blue mackerel catch by country, region and fisheries (ton). The captions in the table below are from the left: Fishing year (July to next June), Total, Total of Japan, Russia, China, Northern purse-seine, Set net, firelight mackerel fishing, Mid-area purse-seine fishery, various fisheries catch at southern area.

漁期年 7月～ 翌年 6月	合計	日本 合計	太平洋北・中区					南区 全漁業	
			ロシア	中国	北区 まき網	定置網 等	火光利 用さば まき網 漁業*		
1982	84,023	84,023	-	-	0	1,354	61,917	826	19,927
1983	65,833	65,833	-	-	0	1,721	36,552	2,308	25,252
1984	92,096	92,096	-	-	0	4,667	55,088	5,816	26,525
1985	120,123	120,123	-	-	0	10,287	62,420	17,092	30,325
1986	107,583	107,583	-	-	532	3,925	53,655	18,010	31,460
1987	97,262	97,262	-	-	0	4,097	35,929	10,532	46,704
1988	57,242	57,242	-	-	0	4,579	18,240	12,067	22,356
1989	47,458	47,458	-	-	0	2,082	15,331	8,034	22,011
1990	27,864	27,864	-	-	47	2,070	7,767	5,678	12,302
1991	23,024	23,024	-	-	113	692	7,164	5,070	9,984
1992	56,060	56,060	-	-	10	4,434	11,870	20,284	19,463
1993	65,231	65,231	-	-	0	3,335	19,511	18,327	24,058
1994	71,962	71,962	-	-	0	2,348	18,718	26,894	24,002
1995	131,067	131,067	-	-	14,824	3,040	21,057	64,498	27,647
1996	179,832	179,832	-	-	13,184	10,938	26,514	72,788	56,408
1997	146,324	146,324	-	-	6,589	5,008	24,871	63,903	45,953
1998	58,385	58,385	-	-	7,641	1,334	15,348	23,544	10,518
1999	121,315	121,315	-	-	14,238	4,381	19,607	56,695	26,393
2000	113,597	113,597	-	-	25,548	3,830	23,365	44,230	16,624
2001	116,056	116,056	-	-	46,230	4,022	18,847	33,817	13,140
2002	110,135	110,135	-	-	11,746	7,802	16,760	46,575	27,252
2003	110,413	110,413	-	-	11,464	5,686	19,948	53,951	19,365
2004	158,927	158,927	-	-	16,673	23,107	18,631	74,934	25,582
2005	191,870	191,870	-	-	35,965	23,182	12,705	73,986	46,032
2006	192,976	192,976	-	-	42,643	20,777	11,890	89,427	28,239
2007	122,171	122,171	-	-	42,627	10,319	13,579	42,525	13,121
2008	149,584	149,584	-	-	46,848	19,624	12,572	45,411	25,129
2009	179,244	179,244	-	-	64,200	13,488	10,643	62,853	28,060
2010	190,993	190,993	-	-	44,136	32,121	13,732	68,058	32,947
2011	180,014	180,014	-	-	54,986	13,537	11,676	66,234	33,580
2012	135,075	135,075	-	-	35,991	14,278	8,015	56,504	20,288
2013	109,998	109,998	-	-	33,088	16,855	7,545	30,294	22,216
2014	120,382	115,192	9	5,180	41,393	17,005	7,568	33,608	15,619
2015	81,431	68,925	43	12,463	15,565	20,473	5,597	18,155	9,134
2016	49,215	46,892	141	2,182	2,907	14,998	7,048	12,500	9,440
2017	36,658	34,345	594	1,720	2,392	11,877	4,584	4,972	10,520
2018	34,879	33,862	439	579	490	8,139	3,640	7,793	13,801

**Table4-1.** Results of the cohort-analysis. The captions in the table are from the left: fishing year, catch (thousand tons), biomass (thousand tons), SSB (thousand tons), recruitment (million fish), catch proportion (%), and RPS (numbers/kg).

漁期年	漁獲量 (千トン)	資源量 (千トン)	親魚量 (千トン)	加入量 (百万尾)	漁獲割合 (%)	再生産成功率 (尾/kg)
1995	131	309	49	732	42	15.0
1996	177	377	38	1,549	47	41.2
1997	149	268	52	528	56	10.1
1998	58	254	50	548	23	11.0
1999	121	363	73	826	33	11.3
2000	114	345	100	649	33	6.5
2001	116	378	104	733	31	7.0
2002	110	342	100	774	32	7.8
2003	109	299	94	825	37	8.8
2004	159	646	81	3,079	25	37.9
2005	192	659	123	858	29	7.0
2006	194	589	340	691	33	2.0
2007	122	587	243	1,166	21	4.8
2008	150	481	197	676	31	3.4
2009	179	720	217	2,314	25	10.7
2010	189	704	168	778	27	4.6
2011	181	591	319	893	31	2.8
2012	137	489	220	720	28	3.3
2013	109	414	200	606	26	3.0
2014	120	316	165	654	38	4.0
2015	83	225	111	204	37	1.8
2016	49	146	83	297	34	3.6
2017	37	113	48	187	33	3.9
2018	34	124	53	292	28	5.6

**Table 4-2.** The abundance indices by various survey. r: regression coefficient between recruitment and SSB. No.1, 2, 5, and 6 indicate regression coefficients using recruitment of beginning year to 2018. No.3 is regression coefficient of recruitment during 2010 to 2018. No. 4 and 4' are it of 2007 to 2019. No.3 and 4' are tuning index of cohort analysis.

	①	②	③	④	④'	⑤	⑥
1995			2,235				5.00
1996			13,870				53.57
1997			321				4.17
1998			8				4.76
1999			2,560				20.83
2000			109				30.77
2001	21.98	14.08	2,142			0.04	24.14
2002	26.98	16.12	813			8.11	36.67
2003	21.52	14.67	692			1.01	25.81
2004	52.73	19.90	30,269			11.11	69.23
2005	29.79	14.69	1,146	4.35	0.48	0.62	3.70
2006	8.70	12.27	59	8.52	0.86	0.01	3.45
2007	36.36	17.70	6,038	53.25	1.89	5.34	30.00
2008	28.85	12.45	716	43.89	2.56	2.13	27.27
2009	42.11	20.14	18,386	16.27	1.76	16.59	58.33
2010	26.32	17.72	3,774	25.56	1.88	1.48	21.74
2011	29.55	14.32	2,939	50.79	1.15	0.22	23.08
2012	35.56	18.41	5,700	26.37	1.81	1.97	30.77
2013	38.64	17.33	5,423	43.12	1.45	104.14	73.68
2014	23.64	17.98	5,718	25.98	1.26	0.14	50.00
2015	29.73	17.86	892	8.87	0.43	2.98	18.18
2016	37.14	18.88	2,505	11.98	0.30	15.00	33.33
2017	53.57	19.86	35	17.27	0.76	21.57	33.33
2018	68.18	18.42	296	109.59	0.39	601.11	69.23
2019	12.50	18.44		10.87	0.14	0.02	19.23
r	0.40	0.02	0.71	0.00	0.55	-0.04	0.40

- ① CPUE of YOY caught by midwater trawl from the spring survey in Northwestern Pacific (number of fish / net / 60 min).
- ② Average length of age 0 fish caught by midwater trawl from the fall survey in Northwestern Pacific
- ③ CPUE of dip-net fishery at the Izu Islands waters (kg / person / hour).
- ④ Number of eggs in region I – III (entire Pacific coast, by 100 billion).
- ④' Standardized number of eggs spawn (Appendix 3).
- ⑤ Estimated abundance of age 0 fish in number (billion fish) from midwater trawl survey in Northwestern Pacific.
- ⑥ Occurrence ration (%) of midwater trawl survey at Northwestern Pacific and drift net survey at offshore Tohoku area.